

Evans-Kishimoto type argument for actions of discrete amenable groups on McDuff factors

MASUDA Toshihiko

Graduate School of Mathematics, Kyushu University,
Fukuoka, 812-5250, JAPAN

Abstract

We apply the Evans-Kishimoto type argument for centrally free actions of discrete amenable groups on McDuff factors, and classify them. Especially, we present a different proof that the Connes-Takesaki modules are complete cocycle conjugacy invariants for centrally free actions of discrete amenable groups on injective factors.

1 Introduction

In the theory of operator algebras, the study of automorphism groups is one of the most important subjects. Especially, since Connes succeeded in classifying automorphisms of the approximately finite dimensional (AFD) factor of type II_1 in [4] and [1], classification of actions of discrete amenable groups on injective factors has been solved in [9], [16], [19], [11] and finally in [10].

The idea of Connes's classification is following. First he constructed tensor product type model automorphisms (or actions) on the AFD type II_1 factor. Then he showed "the model action splitting", i.e., every automorphism contains model actions as tensor product components after an appropriate inner perturbation, and then proved that it is cocycle conjugate to the model action. In his arguments, the main tool is a Rohlin property for automorphisms, (or actions). In [1], he showed the noncommutative version of Rohlin type theorem for a certain class of automorphisms. By means of the Rohlin type theorem, he proved the stability (or 1-cohomology vanishing theorem) for automorphisms, which is the important step in his classification. Connes's argument has been developed by Jones for finite group case in [9], and Ocneanu for general discrete amenable group case in [16]. Especially, the Rohlin type theorem was extended to the case of discrete amenable groups by Ocneanu, and he proved several cohomology vanishing theorems in [16].

On the other hand, another approach has been made in the study of automorphisms of C^* -algebras. In [6], Evans and Kishimoto developed the intertwining argument for classification of automorphisms with the Rohlin property. In their approach, they compare given two automorphisms directly without using model actions. As a consequence, they can treat a wide class of automorphisms, and obtained classification results. (In C^* -algebra case, arguments based on model actions forces us to make a strict restriction

on actions.) Their intertwining argument has been farther developed in [15] for automorphisms of purely infinite simple C^* -algebras, and also for finite group actions in [8].

In this paper, we apply the Evans-Kishimoto type intertwining argument for actions of discrete amenable groups on McDuff factors based on Ocneanu's Rohlin type theorem. Our main theorem says that if two centrally free actions of a discrete amenable group on a McDuff factor differ up to approximately inner automorphisms, then they are cocycle conjugate. Especially if we apply our result to injective factors, then we get the complete classification of centrally free actions of discrete amenable groups in terms of Connes-Takesaki module by using the characterization of approximately inner automorphisms for injective factors in [11]. Hence this is another proof of classification results in [1], [9], [16], [19], [11] for centrally free actions. However our approach seems to be more unified and simple one, and this is an advantage of our theory.

Our result is also applicable to the classification of group actions on subfactors by suitable modification. For examples, we present a different proof of Popa's result in [17, Theorem 3.1]. (We remark that the classification result of strongly amenable subfactors of type II_1 by Popa in [17] is crucial in our argument.)

Acknowledgements. The author is grateful to Professor Izumi, Professor Katayama, and Professor Takesaki for comments on this work. He is supported by Grant-Aid for Scientific Research, Japan Society for the Promotion of Science.

2 Preliminaries and notations

Let M be a von Neumann algebra. For $\varphi \in M_*^+$, we let $\|x\|_\varphi = \sqrt{\varphi(x^*x)}$, $\|x\|_\varphi^\# = \sqrt{(\|x\|_\varphi^2 + \|x^*\|_\varphi^2)/2}$, $|x|_\varphi = \varphi(|x|)$. Note $|x|_\varphi$ is not necessary a norm unless φ is tracial, since it is not subadditive. For $x \in M$ and $\varphi \in M_*$, functionals $x\varphi$ and φx are defined as $x\varphi(y) = \varphi(yx)$ and $\varphi x(y) = \varphi(xy)$ respectively. We let $[x, \varphi] = x\varphi - \varphi x$. To avoid possible confusions, we also denote $x\varphi$ and φx by $x \cdot \varphi$ and $\varphi \cdot x$. We denote $\phi \circ \alpha^{-1}$ by $\alpha(\phi)$ for $\phi \in M_*$ and $\alpha \in \text{Aut}(M)$.

We use the notation $A \subset\subset B$ if A is a finite subset of B , and denote the cardinality of A by $|A|$.

Fix a free ultrafilter ω over $\mathbf{N}$. We define M_ω and M^ω as in [16]. Each $\alpha \in \text{Aut}(M)$ gives an automorphism $\alpha^\omega \in \text{Aut}(M^\omega)$, and $\alpha^\omega|_{M_\omega} \in \text{Aut}(M_\omega)$. For $\varphi \in M_*$ and $X = (x_n) \in M^\omega$, $\varphi^\omega(X) := \lim_{n \rightarrow \omega} \varphi(x_n)$ is a normal functional of M^ω , which we denote by φ for simplicity. When M is a factor, $\tau_\omega(X) := \lim_{n \rightarrow \omega} x_n (\in \mathbf{C})$ always exists in σ -weak topology for $X = (x_n) \in M_\omega$, and τ_ω is a tracial state on M_ω . We denote by $|X|_1$ the L^1 -norm with respect to this trace.

We collect fundamental, frequently used inequalities in this paper.

Lemma 1 *We have the followings for $\varphi \in M_*^+$, $x \in M$.*

- (1) $\|x \cdot \varphi\| \leq \sqrt{\|\varphi\|} \|x\|_\varphi$, $\|\varphi \cdot x\| \leq \sqrt{\|\varphi\|} \|x^*\|_\varphi$, $\|[x, \varphi]\| \leq 2\sqrt{\|\varphi\|} \|x\|_\varphi^\#$.
- (2) $\|x\|_\varphi^2 \leq \|x \cdot \varphi\| \|x\|$, $\|x^*\|_\varphi^2 \leq \|\varphi \cdot x\| \|x\|$.
- (3) $\|x\|_\varphi^\# \leq \sqrt{\frac{1}{2}(|x|_\varphi + |x^*|_\varphi)} \|x\|$.
- (4) For $x_i \in M^\omega$ and $y_i \in M_\omega$, $|\sum_i x_i y_i|_\varphi \leq \sum_i \|x_i\| |y_i|_1$.

Proof. It is elementary to see (1), (2), (3). See [16, Lemma 7.1] for the proof of (4). $\square$

Next we recall Ocneanu's Rohlin type theorem, which is a main tool in the proof of Lemma 6 below.

Theorem 2 ([16, Theorem 6.1]) *Let M be a McDuff factor, G a discrete amenable group, α an action of G on M_ω which is strongly free and semiliftable. Let $\varepsilon > 0$ and let $\{K_i\}_{i \in I}$ be an ε -paving family. Then there exists a partition of unity $\{E_{i,k}\}_{i \in I, k, l \in K_i} \subset M_\omega$ such that*

$$\sum_{i \in I} |K_i|^{-1} \sum_{k \in K_i} |\alpha_{kl^{-1}}(E_{i,l}) - E_{i,k}|_1 \leq 5\varepsilon^{\frac{1}{2}},$$

$$[\alpha_g(E_{i,k}), E_{j,l}] = 0, \text{ for all } g \in G, i, j \in I, k \in K_i, l \in K_j.$$

See [16] for terms appearing in the above theorem. Here we briefly explain how to construct an ε -paving family $\{K_i\}$. Fix $N \in \mathbf{N}$ such that $N > \frac{4}{\varepsilon} \log \varepsilon^{-1}$, and set $\delta := (\varepsilon/3)^N$. Let K_{n+1} be a $(\delta|\bar{K}_n|^{-1}, \bar{K}_n)$ invariant finite set, where $\bar{K}_n := \cup_{1 \leq i \leq n} K_i$. (In this paper, we say K is (ε, F) -invariant if $|K \cap \bigcap_{g \in F} g^{-1}K| \geq (1-\varepsilon)|K|$.) Then $\{K_i\}_{1 \leq i \leq N}$ is shown to be an ε -paving family. In this construction, each K_i can be chosen arbitrarily invariant.

3 Classification

We state the main result in this paper.

Theorem 3 *Let M be a McDuff factor, G a countable discrete amenable group, α, β centrally free actions of G on M . If $\alpha_g \beta_g^{-1} \in \overline{\text{Int}}(M)$ for every $g \in G$, then there exist an α -cocycle v_g and $\theta \in \overline{\text{Int}}(M)$ such that $\text{Adv}_g \alpha_g = \theta \circ \beta_g \circ \theta^{-1}$. Moreover we can choose v_g close to 1, i.e., for given any $\varepsilon > 0$, $F \subset\subset G$ and $\varphi \in M_*^+$, we can choose v_g so that $\|v_g - 1\|_\varphi < \varepsilon$ for any $g \in F$.*

The assumption $\alpha_g \beta_g^{-1} \in \overline{\text{Int}}(M)$ implies that we can approximate α_g by the inner perturbation of β_g . However the inner perturbation of β_g is not necessary an action of G . Hence our first task is to approximate α by the cocycle perturbation of β .

Proposition 4 *Under the assumption in Theorem 3, there exist β -cocycles u_g^n , $n = 1, 2, \dots$ such that $\alpha_g = \lim_{n \rightarrow \omega} \text{Ad} u_g^n \beta_g$.*

Proof. Since $\alpha_g \beta_g^{-1} \in \overline{\text{Int}}(M)$, there exist unitaries u_g^n , $g \in G$, $n = 1, 2, \dots$ such that $\alpha_g = \lim_{n \rightarrow \omega} \text{Ad} u_g^n \beta_g$. Set $U_g := (u_g^n) \in M^\omega$. Then $\alpha_g^\omega = \text{Ad} U_g \beta_g^\omega$ holds on $M(\subset M^\omega)$. Set $u(g, h) = U_g \beta_g^\omega(U_h) U_{gh}^*$. Then it is easy to check $u(g, h) \in M_\omega$ and $\{\text{Ad} U_g \beta_g^\omega|_{M_\omega}, u(g, h)\}$ is a cocycle action on M_ω . Moreover $\text{Ad} U_g \beta_g^\omega$ is strongly free in the sense of [16, Definition 5.6] by [16, Lemma 5.7]. By Ocneanu's 2-cohomology vanishing theorem [16, Proposition 7.4], we get $c_g \in U(M_\omega)$ such that $c_g \text{Ad} U_g \beta_g^\omega(c_h) u(g, h) c_{gh}^* = 1$. Let $c_g = (c_g^n)$ be a representing sequence consisting of unitaries. Set $W_g := c_g U_g = (c_g^n u_g^n)$. Since (c_g^n) is a centralising sequence, $\text{Ad} c_g^n$ converges to id_M . Hence $\alpha_g = \lim_{n \rightarrow \omega} \text{Ad} c_g^n u_g^n \beta_g$, and W_g is a β^ω -cocycle. Set $w_g^n = c_g^n u_g^n$ and $w'_n(g, h) := w_g^n \beta_g(w_h^n) w_{gh}^{n*}$. Then $(\text{Ad} w_g^n \beta_g, w'_n(g, h))$ is a cocycle action, and the cocycle identity $W_g \beta_g(W_h) = W_{gh}$ implies $\lim_{n \rightarrow \omega} w'_n(g, h) = 1$ in the σ -strong* topology. By Ocneanu's 2-cohomology vanishing theorem [16, Theorem 7.6], we can find $d_g^n \in U(M)$ such that $d_g^n \text{Ad} w_g^n \beta_g(d_h^n) w'_n(g, h) d_{gh}^{n*} = 1$ and $\lim_{n \rightarrow \omega} d_g^n = 1$

in the σ -strong* topology. Then $d_g^n w_g^n$ is a β_g -cocycle and $(d_g^n w_g^n) = (w_g^n)$ in M^ω . It is easy to see $\alpha_g \beta_g^{-1} = \lim_{n \rightarrow \omega} \text{Add}_g^n w_g^n$. $\square$.

We get the following corollary immediately.

Corollary 5 *Let α, β be as in Theorem 3. Then for any $\varepsilon > 0$, $\Phi \subset\subset M_*$ and $F \subset\subset G$, there exists a β -cocycle u_g such that $\|\alpha_g(\phi) - \text{Adu}_g \beta_g(\phi)\| < \varepsilon$ for every $g \in F$ and $\phi \in \Phi$.*

Next we show an almost 1-cohomology vanishing, which is essential in our argument.

Lemma 6 *Let α be a centrally free action of G on M . For any $\varepsilon > 0$, $F = F^{-1} \subset\subset G$, $\Phi^+ \subset\subset M_*^+$, and $\Phi \subset\subset M_*$, there exist $\delta > 0$ and $\Psi \subset\subset M_*$ with the following property; for any α -cocycle $\{v_g\}$ with $\|[v_g, \psi]\| < \delta$, $g \in F$, $\psi \in \Psi$, we can find $w \in U(M)$ such that $\|[w, \varphi]\| < \varepsilon$ for every $\varphi \in \Phi$, and $\|v_g \alpha_g(w^*)w - 1\|_\phi^\# < \varepsilon$ for every $g \in F$ and $\phi \in \Phi^+$. When $\Phi = \emptyset$, then $\Psi = \emptyset$ is possible.*

Proof. Since every $\phi \in M_*$ is decomposed as $\phi = \phi_1 - \phi_2 + i(\phi_3 - \phi_4)$, $\phi_i \in M_*^+$, it is sufficient to show the lemma in the case $\Phi = \Phi^+ \subset\subset M_*^+$. We may assume $0 < \varepsilon < 1$ and $\|\phi\| \leq 1$, $\phi \in \Phi^+$. Fix $\varepsilon' > 0$ with $\varepsilon' < (\varepsilon/8)^4$. Let $\{K_i\}_{i \in I}$ be an ε' -paving family such that each K_i is (ε', F) invariant. We may assume that K_i 's are in a subgroup of G generated by F . Define $\text{Length}(g) := \min\{n \mid g = h_1 h_2 \cdots h_n, h_i \in F\}$, and define L by

$$L := \max\{\text{Length}(g) \mid g \in K_i, i \in I\}.$$

Fix $\delta > 0$ so that $\sum_i |K_i|(L+1)\delta < \varepsilon/3$. Define Ψ by

$$\Psi := \Phi^+ \cup \bigcup_{\substack{1 \leq k \leq L-1, \\ g_i \in F}} \alpha_{g_1 g_2 \cdots g_k}^{-1}(\Phi^+).$$

By Ocneanu's Rohlin Theorem there exists a partition of unity $\{E_{i,k}\}_{i \in I, k \in K_i} \subset M_\omega$ such that

$$\sum_{i \in I} |K_i|^{-1} \sum_{k, l \in K_i} |\alpha_{kl^{-1}}(E_{i,l}) - E_{i,k}|_1 < 5\varepsilon'^{\frac{1}{2}},$$

$$[\alpha_g(E_{i,k}), E_{j,l}] = 0 \text{ for all } g \in G, i, j \in I, k \in K_i, l \in K_j.$$

Then by [16, Corollary 6.1], we have followings.

$$\sum_{i \in I} \sum_{k \in K_i \cap g^{-1}K_i} |\alpha(E_{i,k}) - E_{i,gk}|_1 \leq 10\varepsilon'^{\frac{1}{2}} \text{ for every } g \in F,$$

$$\sum_{i \in I} \sum_{k \in A_i} |E_{i,k}|_1 \leq \delta' + 5\varepsilon'^{\frac{1}{2}},$$

for any subsets $A_i \subset K_i$ with $|A_i| \leq \delta' |K_i|$, $i \in I$.

Let v_g be an α -cocycle with $\|[v_g, \phi]\| < \delta$, $g \in F$, $\phi \in \Psi$. Define $W := \sum_{i,k} v_k^* E_{i,k}$. First we estimate $\|v_g \alpha_g(W^*)W - 1\|_\phi^\#$ as in the proof of [16, Proposition 7.2]. We investigate $|v_g \alpha_g(W^*)W - 1|_\phi$ and $|(v_g \alpha_g(W^*)W - 1)^*|_\phi$, then use Lemma 1.

We divide $v_g \alpha_g(W^*)W - 1 = \sum_{i \in I, k \in K_i} \sum_{j \in I, l \in K_j} (v_g \alpha_g(v_k) v_l^* - 1) \alpha_g(E_{i,k}) E_{j,l}$ into three parts as follows.

$$\begin{aligned}
\sum_{i \in I, k \in K_i} \sum_{j \in I, l \in K_j} (*) &= \sum_{j \in I, l \in K_j} \sum_{i, k \in K_i \cap g^{-1} K_i} (*) + \sum_{j \in I, l \in K_j} \sum_{i \in I, k \in K_i \setminus g^{-1} K_i} (*) \\
&= \sum_{j \in I, l \in K_j} \sum_{\substack{j \neq i \in I, \\ k \in K_i \cap g^{-1} K_i}} (*) + \sum_{\substack{j \in I, l \in K_j, \\ k \in K_j \cap g^{-1} K_j}} (*) + \sum_{j \in I, l \in K_j} \sum_{i, k \in K_i \setminus g^{-1} K_i} (*) \\
&= \sum_{\substack{j \in I, l \in K_j, \\ k \in K_j \cap g^{-1} K_j, gk=l}} (*) + \sum_{\substack{j \in I, l \in K_j, \\ k \in K_j \cap g^{-1} K_j, gk \neq l}} (*) \\
&\quad + \sum_{j \in I, l \in K_j} \sum_{\substack{j \neq i \in I, \\ k \in K_i \cap g^{-1} K_i}} (*) + \sum_{j \in I, l \in K_j} \sum_{i, k \in K_i \setminus g^{-1} K_i} (*) \\
&= \sum_1 (*) + \sum_2 (*) + \sum_3 (*).
\end{aligned}$$

In $\sum_1$ we sum for $i = j, k \in K_i \cap g^{-1} K_i, gk = l$, in $\sum_2$ we sum for $i = j, k \in K_i \cap g^{-1} K_i, gk \neq l$, or $i \neq j, k \in K_i \cap g^{-1} K_i, l \in K_j$, in $\sum_3$ we sum for $k \in K_i \setminus g^{-1} K_i$. Due to the cocycle identity, we have $\sum_1 (v_g \alpha_g(v_k) v_l^* - 1) \alpha_g(E_{i,k}) E_{j,l} = \sum_j (v_g \alpha_g(v_k) v_{gk}^* - 1) \alpha_g(E_{i,k}) E_{j,l} = 0$, and $\sum_1$ part vanishes. Hence

$$\begin{aligned}
&|v_g \alpha_g(W^*)W - 1|_\phi \\
&= \left| \sum_2 (v_g \alpha_g(v_k) v_l^* - 1) \alpha_g(E_{i,k}) E_{j,l} + \sum_3 (v_g \alpha_g(v_k) v_l^* - 1) \alpha_g(E_{i,k}) E_{j,l} \right|_\phi \\
&\leq 2 \sum_2 |\alpha_g(E_{i,k}) E_{j,l}|_1 + 2 \sum_3 |\alpha_g(E_{i,k}) E_{j,l}|_1
\end{aligned}$$

holds by Lemma 1(4). Similarly we have

$$|(v_g \alpha_g(W^*)W - 1)^*|_\phi \leq 2 \sum_2 |\alpha_g(E_{i,k}) E_{j,l}|_1 + 2 \sum_3 |\alpha_g(E_{i,k}) E_{j,l}|_1.$$

The estimation of $\sum_3$ is as follows.

$$\begin{aligned}
\sum_3 |\alpha_g(E_{i,k}) E_{j,l}|_1 &= \sum_{j \in I, l \in K_j} \sum_{\substack{i \in I, \\ k \in K_i \setminus g^{-1} K_j}} |\alpha_g(E_{i,k}) E_{j,l}|_1 \\
&= \sum_{\substack{i \in I, \\ k \in K_j \setminus g^{-1} K_j}} |E_{i,k}|_1 \\
&\leq \varepsilon' + 5\varepsilon'^{\frac{1}{2}} \leq 6\varepsilon'^{\frac{1}{2}},
\end{aligned}$$

since we have $|K_j \setminus g^{-1} K_j| \leq \varepsilon' |K_j|$.

Next we estimate $\sum_2$ part. Then

$$\begin{aligned}
\sum_2 |\alpha_g(E_{i,k}) E_{j,l}|_1 &= \sum_{\substack{j \in I, \\ l \in K_j}} \sum_{\substack{i \neq j, \\ k \in K_i \cap g^{-1} K_i}} |\alpha_g(E_{i,k}) E_{j,l}|_1 + \sum_{\substack{j \in I, l \in K_j, \\ k \in K_j \cap g^{-1} K_j, gk \neq l}} |\alpha_g(E_{i,k}) E_{j,l}|_1 \\
&= \sum_{\substack{i \in I, \\ k \in K_i \cap g^{-1} K_i}} |\alpha(E_{i,k}) (1 - \sum_l E_{i,l})|_1 + \sum_{\substack{i \in I, \\ k \in K_i \cap g^{-1} K_i}} |\alpha_g(E_{i,k}) \sum_{l \in K_i, l \neq gk} E_{i,l}|_1
\end{aligned}$$

$$\begin{aligned}
&= \sum_{i \in I, k \in K_i \cap g^{-1}K_i} |\alpha_g(E_{i,k})(1 - E_{i,gk})|_1 \\
&= \sum_{i \in I, k \in K_i \cap g^{-1}K_i} |(\alpha_g(E_{i,k}) - E_{i,gk})(1 - E_{i,gk})|_1 \\
&\leq \sum_{i \in I, k \in K_i \cap g^{-1}K_i} |(\alpha_g(E_{i,k}) - E_{i,gk})|_1 \\
&\leq 10\varepsilon'^{\frac{1}{2}}
\end{aligned}$$

holds. Then we get $|v_g \alpha_g(W^*)W - 1|_\phi \leq 32\varepsilon'^{\frac{1}{2}}$ and $|(v_g \alpha_g(W^*)W - 1)^*|_\phi \leq 32\varepsilon'^{\frac{1}{2}}$. By Lemma 1 (3), we have

$$\|v_g \alpha_g(W^*)W - 1\|_\phi^\# \leq 8\varepsilon'^{\frac{1}{4}} < \varepsilon.$$

We choose a representing sequence $W = (w_n)$ consisting of unitaries, and $E_{i,k} = (e_{i,k}^n)$ such that $\{e_{i,k}^n\}_{i \in I, k \in K_i}$ is a partition of unity for each n . Set $a_n := \sum_{i,k} v_k^* e_{i,k}^n$. (Note that a_n is not necessary a unitary.) Since $W = (w_n) = (a_n)$ in M^ω , $\{w_n - a_n\}$ converges to 0 σ -strongly*. Choose a sufficiently large n such that

$$\|v_g \alpha_g(w_n^*)w_n - 1\|_\phi^\# < \varepsilon, g \in F, \phi \in \Phi^+,$$

$$\|[\phi, e_{i,k}^n]\| < \delta, \phi \in \Phi^+, i \in I, k \in K_i,$$

$$\|w_n - a_n\|_\phi^\# < \varepsilon/3, \phi \in \Phi^+.$$

Set $w := w_n$, $a := a_n$, $e_{i,k} := e_{i,k}^n$. (Note that we never use the approximate commutativity of v_g with elements in Ψ in the estimate of $\|v_g \alpha_g(w^*)w - 1\|_\phi^\#$.) Next we estimate $\|[a, \varphi]\|$, $\varphi \in \Phi^+$. To do so, we estimate $\|[v_g, \varphi]\|$, $g \in K_i$, $\varphi \in \Phi^+$ at first. We express g as $g = g_1 g_2 \cdots g_k$, $g_i \in F$, $k \leq L$. Then it is easily shown $\|[v_g, \varphi]\| \leq \sum_{i=1}^k \|[v_{g_i}, \alpha_{g_1 \cdots g_{i-1}}^{-1}(\varphi)]\|$. (When $i = 1$, $\alpha_{g_1 \cdots g_0}(\varphi)$ means φ .) By the definition of Ψ , each $\alpha_{g_1 \cdots g_i}^{-1}(\varphi)$ is in Ψ , and by the assumption on v_g , $\|[v_{g_i}, \alpha_{g_1 \cdots g_{i-1}}^{-1}(\varphi)]\| < \delta$ follows. Hence we have $\|[v_g, \varphi]\| \leq L\delta$. Finally,

$$\begin{aligned}
\|[a, \varphi]\| &\leq \sum_{i \in I, k \in K_i} \|[v_k^* e_{i,k}, \varphi]\| \\
&\leq \sum_{i \in I, k \in K_i} \|[v_k^*, \varphi]e_{i,k}\| + \|v_k^*[e_{i,k}, \varphi]\| \\
&\leq \sum_{i \in I, k \in K_i} (L+1)\delta \\
&= \sum_{i \in I} |K_i|(L+1)\delta \\
&< \varepsilon/3
\end{aligned}$$

holds for $\varphi \in \Phi^+$.

By Lemma 1 (1),

$$\begin{aligned}
\|[w, \varphi]\| &\leq \|[a, \varphi]\| + \|[w - a, \varphi]\| \\
&< \varepsilon/3 + 2\|w - a\|_\phi^\# \\
&< \varepsilon
\end{aligned}$$

holds for $\varphi \in \Phi^+$, and w is a desired unitary. $\square$

Remark. If we replace $v_g \alpha_g(W^*)W$ with $W v_g \alpha_g(W^*)$ in the above proof, then we get a similar result as in Lemma 6 by replacing $\|v_g \alpha_g(w^*)w - 1\|_\phi^\#$ with $\|w v_g \alpha_g(w^*) - 1\|_\phi^\#$.

Now we can present a proof of Theorem 3 by means of Lemma 6.

Proof of Theorem 3. Fix a faithful normal state φ_0 . Let $\Phi = \{\varphi_i\}_{i=0}^\infty$ be a countable dense subset of M_* , and set $\Phi_n := \{\varphi_i\}_{i=0}^n$. Let $G_n \subset\subset G$ be such that $G_n \subset G_{n+1}$, $G_n^{-1} = G_n$ and $\cup_n G_n = G$.

We construct $w_n, v_g^n, \bar{v}_g^n \in U(M)$, $\Phi'_n, \Psi_n \subset\subset M_*$, $\Phi_n^+ \subset\subset M_*^+$, $\delta_n > 0$, actions $\alpha_g^{(2n)}, \beta_g^{(2n-1)}$ of G satisfying the below conditions inductively. (We set $\alpha^{(0)} := \alpha$, $\beta^{(-1)} := \beta$.)

$$(1.2n) \quad \|\beta_g^{(2n-1)}(\varphi) - \alpha_g^{(2n)}(\varphi)\| < 1/2^{2n}, g \in G_{2n}, \varphi \in \Phi'_{2n}, (n \geq 1).$$

$$(1.2n+1) \quad \|\alpha_g^{(2n)}(\varphi) - \beta_g^{(2n+1)}(\varphi)\| < 1/2^{2n+1}, g \in G_{2n+1}, \varphi \in \Phi'_{2n+1}, (n \geq 0).$$

$$(2.2n) \quad \|\beta_g^{(2n-1)}(\psi) - \alpha_g^{(2n)}(\psi)\| < \frac{\delta_{2n-1}}{2}, g \in G_{2n-1}, \psi \in \bigcup_{g \in G_{2n-1}} \beta_{g^{-1}}^{(2n-1)}(\Psi_{2n-1}), (n \geq 1).$$

$$(2.2n+1) \quad \|\alpha_g^{(2n)}(\psi) - \beta_g^{(2n+1)}(\psi)\| < \frac{\delta_{2n}}{2}, g \in G_{2n}, \psi \in \bigcup_{g \in G_{2n}} \alpha_{g^{-1}}^{(2n)}(\Psi_{2n}), (n \geq 1).$$

$$(3.n) \quad \|v_g^n - 1\|_\phi^\# < 1/4^n, g \in G_{n-2}, \phi \in \Phi_{n-2}^+, (G_{-1} = G_0 = G_1, \Phi_{-1}^+ = \Phi_0^+ = \{\varphi_0\}),$$

$$(4.n) \quad \|[w_n, \varphi]\| < 1/4^n, \varphi \in \Phi'_{n-1}, (n \geq 3).$$

$$\bar{v}_g^n := v_g^n \text{Ad} w_n (\bar{v}_g^{n-2}), (\bar{v}_g^1 = v_g^1, \bar{v}_g^2 = v_g^2),$$

$$\alpha_g^{(2n)} := \text{Ad} v_g^{2n} \circ \text{Ad} w_{2n}^* \circ \alpha_g^{(2n-2)} \circ \text{Ad} w_{2n},$$

$$\beta_g^{(2n-1)} := \text{Ad} v_g^{2n-1} \circ \text{Ad} w_{2n-1}^* \circ \beta_g^{(2n-3)} \circ \text{Ad} w_{2n-1},$$

$$\Phi'_{2n} := \Phi_{2n} \cup \text{Ad} w_{2n-1}^* w_{2n-3}^* \cdots w_1^*(\Phi_{2n}) \cup \{\bar{v}_g^{2n-1} \varphi_0, \varphi_0 \bar{v}_g^{2n-1}\}_{g \in G_{2n-1}},$$

$$\Phi'_{2n+1} := \Phi_{2n+1} \cup \text{Ad} w_{2n}^* w_{2n-2}^* \cdots w_2^*(\Phi_{2n+1}) \cup \{\bar{v}_g^{2n} \varphi_0, \varphi_0 \bar{v}_g^{2n}\}_{g \in G_{2n}}, (n \geq 1)$$

$$\Phi_n^+ := \{\text{Ad} \bar{v}_g^n(\varphi_0) \mid g \in G_n\}.$$

Here δ_{2n} and Ψ_{2n} (δ_{2n-1} and Ψ_{2n-1}) are chosen as in Lemma 6 for $\alpha^{(2n)}$, $1/4^{2n+2}$, G_{2n} , Φ_{2n}^+ and Φ'_{2n+1} (resp. for $\beta^{(2n-1)}$, $1/4^{2n+1}$, G_{2n-1} , Φ_{2n-1}^+ , and Φ'_{2n}).

First set $\Phi'_1 := \Phi_1$ and fix a β -cocycle u_g^1 such that

$$\|\alpha_g^{(0)}(\varphi) - \text{Ad} u_g^1 \beta_g^{(-1)}(\varphi)\| < 1/2, g \in G_1, \varphi \in \Phi'_1.$$

By Lemma 6, we get a unitary w_1 such that $\|u_g \beta_g^{(-1)}(w_1^*)w_1 - 1\|_{\varphi_0}^\# < 1/4$, $g \in G_1$. Set $v_g^1 := u_g^1 \beta_g^{(-1)}(w_1^*)w_1$, and $\beta_g^{(1)} := \text{Ad} u_g^1 \circ \beta_g^{(-1)} = \text{Ad} v_g^1 \circ \text{Ad} w_1^* \circ \beta_g^{(-1)} \circ \text{Ad} w_1$. Then we have

$$(1.1) \quad \|\alpha_g^{(0)}(\varphi) - \beta_g^{(1)}(\varphi)\| < 1/2$$

and

$$(3.1) \quad \|v_g^1 - 1\|_{\varphi_0}^\# < 1/4$$

for $g \in G_1$. Set $\bar{v}_g^1 := v_g^1$, $\Phi'_2 := \Phi_2 \cup \text{Ad} w_1^*(F_2) \cup \{\bar{v}_g^1 \varphi_0, \varphi_0 \bar{v}_g^1\}$, and $\Phi_1^+ := \{\text{Ad} \bar{v}_g^1(\varphi_0)\}_{g \in G_1}$. By Lemma 6, we choose Ψ_1 and δ_1 for $\beta_g^{(1)}$, $1/4^3$, G_1 , Φ_1^+ and Φ'_2 .

Next we take an $\alpha^{(0)}$ -cocycle u_g^2 such that

$$(a.2) \quad \|\beta_g^{(1)}(\varphi) - \text{Ad} u_g^2 \alpha_g^{(0)}(\varphi)\| < \frac{1}{2^2}, g \in G_2, \varphi \in \Phi'_2,$$

$$(b.2) \quad \|\beta_g^{(1)}(\psi) - \text{Ad} u_g^2 \alpha_g^{(0)}(\psi)\| < \frac{\delta_1}{2}, g \in G_1, \psi \in \bigcup_{g \in G_1} \beta_{g^{-1}}^{(1)}(\Psi_1).$$

By Lemma 6, we get $w_2 \in U(M)$ such that $\|u_g^2 \alpha_0(w_2^*) w_2 - 1\|_{\varphi_0}^\# < 1/4^2$ for $g \in G_2$. Set $v_g^2 = \bar{v}_g^2 := u_g^2 \alpha_g^{(0)}(w_2^*) w_2$ and $\alpha_g^{(2)} := \text{Adu}_g^2 \alpha_g^{(0)} = \text{Ad} v_g^2 \circ \text{Ad} w_2^* \circ \alpha_g^{(0)} \circ \text{Ad} w_2$. Then we get

$$(3.2) \quad \|v_g^2 - 1\|_{\varphi_0}^\# < \frac{1}{4^2}, g \in G_2.$$

By (a.2) and (b.2),

$$(1.2) \quad \|\beta_g^{(1)}(\psi) - \alpha_g^{(2)}(\psi)\| < \frac{1}{2^2}, g \in G_2, \varphi \in \Phi'_2,$$

$$(2.2) \quad \|\beta_g^{(1)}(\psi) - \alpha_g^{(2)}(\psi)\| < \frac{\delta_1}{2}, g \in G_1, \psi \in \bigcup_{g \in G_1} \beta_{g^{-1}}^{(1)}(\Psi_1).$$

Set $\Phi'_3 := \Phi_3 \cup \text{Ad} w_2^*(\Phi_3) \cup \{\bar{v}_g^2 \varphi_0, \varphi_0 \bar{v}_g^2\}_{g \in G_2}$ and $\Phi_2^+ := \{\text{Ad} \bar{v}_g^2 \varphi_0 \mid g \in G_2\}$. By Lemma 6, we choose δ_2 and Ψ_2 for $\alpha^{(2)}$, G_2 , Φ'_3 , Φ_2^+ and $1/4^4$.

Suppose that we construct up to $\alpha_g^{(2n)}$, $\beta_g^{(2n-1)}$, w_{2n} , v_g^{2n} , $\bar{v}_g^{2n}$, Φ'_{2n+1} , δ_{2n} and Ψ_{2n} .

We choose a $\beta^{(2n-1)}$ -cocycle $u_g^{2n+1} \in U(M)$ such that

$$(a.2n+1) \quad \|\alpha_g^{(2n)}(\varphi) - \text{Adu}_g^{2n+1} \beta_g^{(2n-1)}(\varphi)\| < \frac{1}{2^{2n+1}}, g \in G_{2n+1}, \varphi \in \Phi'_{2n+1},$$

$$(b.2n+1) \quad \|\alpha_g^{(2n)}(\psi) - \text{Adu}_g^{2n+1} \beta_g^{(2n-1)}(\psi)\| < \frac{\delta_{2n}}{2}, g \in G_{2n}, \psi \in \bigcup_{g \in G_{2n}} \alpha_{g^{-1}}^{(2n)}(\Psi_{2n}),$$

$$(c.2n+1) \quad \|\alpha_g^{(2n)}(\psi) - \text{Adu}_g^{2n+1} \beta_g^{(2n-1)}(\psi)\| < \frac{\delta_{2n-1}}{2}, g \in G_{2n-1}, \varphi \in \bigcup_{g \in G_{2n-1}} \beta_{g^{-1}}^{(2n-1)}(\Psi_{2n-1}).$$

Then by (2.2n) and (c.2n+1) we get

$$\|\beta_g^{(2n-1)}(\psi) - \text{Adu}_g^{2n+1} \beta_g^{(2n-1)}(\psi)\| < \delta_{2n-1}, g \in G_{2n-1}, \psi \in \beta_{g^{-1}}^{(2n-1)}(\Psi_{2n-1}),$$

which yields $\|u_g^{2n+1} \beta_g^{(2n-1)}(\psi) - \beta_g^{(2n-1)}(\psi)\| < \delta_{2n-1}$, $g \in G_{2n-1}$, $\psi \in \Psi_{2n-1}$. By the choice of δ_{2n-1} and Ψ_{2n-1} , there exists a unitary w_{2n+1} such that

$$\|u_g^{2n+1} \beta_g^{(2n-1)}(w_{2n+1}^*) w_{2n+1} - 1\|_{\phi}^\# < 1/4^{2n+1}, g \in G_{2n-1}, \phi \in \Phi_{2n-1}^+,$$

and

$$(4.2n+1) \quad \|[w_{2n+1}, \varphi]\| < 1/4^{2n+1}, \varphi \in \Phi'_{2n}.$$

Set $v_g^{2n+1} := u_g^{2n+1} \beta_g^{(2n-1)}(w_{2n+1}^*) w_{2n+1}$ and $\beta_g^{(2n+1)} := \text{Adu}_g^{(2n+1)} \beta_g^{(2n-1)} = \text{Ad} v_g^{(2n+1)} \circ \text{Ad} w_{2n+1}^* \circ \beta_g^{(2n-1)} \circ \text{Ad} w_{2n+1}$. Then

$$(3.2n+1) \quad \|v_g^{2n+1} - 1\|_{\phi} < \frac{1}{4^{2n+1}}, g \in G_{2n-1}, \phi \in \Phi_{2n-1}^+.$$

By (a.2n+1) and (b.2n+1), we get

$$(1.2n+1) \quad \|\alpha_g^{(2n)}(\varphi) - \beta_g^{(2n+1)}(\varphi)\| < \frac{1}{2^{2n+1}}, g \in G_{2n+1}, \varphi \in \Phi'_{2n+1}.$$

$$(2.2n+1) \quad \|\alpha_g^{(2n)}(\psi) - \beta_g^{(2n+1)}(\psi)\| < \frac{\delta_{2n}}{2}, g \in G_{2n}, \psi \in \bigcup_{g \in G_{2n}} \alpha_{g^{-1}}^{(2n)}(\Psi_{2n}).$$

Set $\bar{v}_g^{2n+1} := v_g^{2n+1} \text{Ad} w_{2n+1}^* (\bar{v}_g^{2n-1})$, and define

$$\Phi'_{2n+2} := \Phi_{2n+2} \cup \text{Ad} w_{2n+1}^* w_{2n-1}^* \cdots w_1^* (\Phi_{2n+2}) \cup \{\bar{v}_g^{2n+1}(\varphi_0), \varphi_0 \bar{v}_g^{2n+1}\}_{g \in G_{2n+1}},$$

$$\Phi_{2n+1}^+ := \{\text{Ad} \bar{v}_g^{2n+1} \varphi_0 \mid g \in G_{2n+1}\}.$$

By Lemma 6, we choose $\delta_{2n+1} > 0$ and $\Psi_{2n+1} \subset\subset M_*$ for $\beta^{(2n+1)}$, $1/4^{2n+3}$, G_{2n+1} , Φ_{2n+1}^+ and Φ'_{2n+2} , and the $2n$ -th step is finished.

Next we choose an $\alpha^{(2n)}$ -cocycle u_g^{2n+2} such that

$$\begin{aligned} (a.2n+2) \quad & \|\beta_g^{(2n+1)}(\varphi) - \text{Ad} u_g^{2n+2} \alpha_g^{(2n)}(\varphi)\| < \frac{1}{2^{2n}}, \quad g \in G_{2n+2}, \varphi \in \Phi'_{2n+2}, \\ (b.2n+2) \quad & \|\beta_g^{(2n+1)}(\psi) - \text{Ad} u_g^{2n+2} \alpha_g^{(2n)}(\psi)\| < \frac{\delta_{2n+1}}{2}, \quad g \in G_{2n+1}, \psi \in \bigcup_{g \in G_{2n+1}} \beta_{g^{-1}}^{(2n+1)}(\Psi_{2n+1}), \\ (c.2n+2) \quad & \|\beta_g^{(2n+1)}(\psi) - \text{Ad} u_g^{2n+2} \alpha_g^{(2n)}(\psi)\| < \frac{\delta_{2n}}{2}, \quad g \in G_{2n}, \psi \in \bigcup_{g \in G_{2n}} \alpha_{g^{-1}}^{(2n)}(\Psi_{2n}). \end{aligned}$$

By (c.2n+2) and (2.2n+1), we get

$$\|\alpha_g^{(2n)}(\psi) - \text{Ad} u_g^{2n+2} \alpha_g^{(2n)}(\psi)\| < \delta_{2n}, \quad g \in G_{2n}, \psi \in \bigcup_{g \in G_{2n}} \alpha_{g^{-1}}^{(2n)}(\Psi_{2n}).$$

We thus have $\|[u_g^{2n+2}, \psi]\| < \delta_{2n}$ for $g \in G_{2n}$ and $\psi \in \Psi_{2n}$. By the choice of δ_{2n} and Ψ_{2n} , we can find $w_{2n+2} \in U(M)$ such that

$$\|u_g^{2n+2} \alpha_g^{(2n)}(w_{2n+2}^*) w_{2n+2} - 1\|_\phi^\# < \frac{1}{4^{2n+2}}, \quad g \in G_{2n}, \phi \in \Phi_{2n}^+$$

and

$$(4.2n+2) \quad \|[w_{2n+2}, \varphi]\| < \frac{1}{4^{2n+2}}, \quad \varphi \in \Phi'_{2n+1}.$$

Set $v_g^{2n+2} := u_g^{2n+2} \alpha_g^{(2n)}(w_{2n+2}^*) w_{2n+2}$ and $\alpha_g^{(2n+2)} := \text{Ad} u_g^{2n+2} \alpha_g^{(2n)} = \text{Ad} v_g^{2n+2} \circ \text{Ad} w_{2n+2}^* \circ \alpha_g^{(2n)} \circ \text{Ad} w_{2n+2}$. Then

$$(3.2n+2) \quad \|v_g^{2n+2} - 1\|_\phi^\# < \frac{1}{4^{2n+2}}, \quad g \in G_{2n}, \phi \in \Phi_{2n}^+,$$

and by (a.2n+2) and (b.2n+2), we get

$$\begin{aligned} (1.2n+2) \quad & \|\beta_g^{(2n+1)}(\varphi) - \alpha_g^{(2n+2)}(\varphi)\| < \frac{1}{2^{2n}}, \quad g \in G_{2n}, \varphi \in \Phi'_{2n+2}. \\ (2.2n+2) \quad & \|\beta_g^{(2n+1)}(\psi) - \alpha_g^{(2n+2)}(\psi)\| < \frac{\delta_{2n+1}}{2}, \quad g \in G_{2n+1}, \psi \in \bigcup_{g \in G_{2n+1}} \beta_{g^{-1}}^{(2n+1)}(\Psi_{2n+1}). \end{aligned}$$

Set $\bar{v}_g^{2n+2} := v_g^{2n+2} \text{Ad} w_{2n+2}^* (\bar{v}_g^{2n})$ and

$$\Phi'_{2n+3} := \Phi_{2n+3} \cup \text{Ad} w_{2n+2}^* w_{2n}^* \cdots w_2^* (\Phi_{2n+3}) \cup \{\bar{v}_g^{2n+2} \varphi_0, \varphi_0 \bar{v}_g^{2n+2}\},$$

$$\Phi_{2n+2}^+ = \{\text{Ad} \bar{v}_g^{2n+2} \varphi_0 \mid g \in G_{2n+2}\}.$$

We choose $\delta_{2n+2} > 0$, $\Psi_{2n+2} \subset\subset M_*$ for $\alpha_g^{(2n)}$, $1/4^{2n+4}$, G_{2n+2} , Φ_{2n+2}^+ and Φ'_{2n+3} by Lemma 6. Then the $(2n+1)$ -th step is finished, and thus we complete induction.

Set $\theta_{2n} := \text{Ad} w_{2n}^* w_{2n-2}^* \cdots w_2^*$. Then we have $\alpha_g^{(2n)} = \text{Ad} \bar{v}_g^{2n} \circ \theta_{2n} \circ \alpha_g \circ \theta_{2n}^{-1}$. We will verify $\{\theta_{2n}\}$ converges to some $\theta \in \text{Aut}(M)$. To do so, we will prove $\{\theta_{2n}(\varphi)\}$ and $\{\theta_{2n}^{-1}(\varphi)\}$ are Cauchy sequences for $\varphi \in M_*$. Suppose $\varphi \in \Phi_k$. For any n with $k \leq 2n+1$, φ and $\theta_{2n}(\varphi)$ are in Φ'_{2n+1} . By $(4.2n+2)$, we have

$$\begin{aligned} \|\theta_{2n+2}(\varphi) - \theta_{2n}(\varphi)\| &= \| [w_{2n+2}, \theta_{2n}(\varphi)] \| \\ &< \frac{1}{4^{2n+2}}, \end{aligned}$$

and

$$\begin{aligned} \|\theta_{2n+2}^{-1}(\varphi) - \theta_{2n}^{-1}(\varphi)\| &= \| w_{2n+2}^* \varphi w_{2n+2} - \varphi \| \\ &< \frac{1}{4^{2n+2}}. \end{aligned}$$

It follows that $\{\theta_{2n}(\varphi)\}$ and $\{\theta_{2n}^{-1}(\varphi)\}$ are Cauchy sequences for $\varphi \in \Phi$. Then so are $\{\theta_{2n}(\varphi)\}$ and $\{\theta_{2n}^{-1}(\varphi)\}$ for every $\varphi \in M_*$, since Φ is dense in M_* . Hence $\{\theta_{2n}\}$ converges to some $\theta \in \text{Aut}(M)$.

Next we will verify $\{\bar{v}_g^{2n}\}$ is a Cauchy sequence with respect to $\|\cdot\|_{\varphi_0}^\#$. Since we have

$$\begin{aligned} \|\bar{v}_g^{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^\# &\leq \| (v_g^{2n+2} - 1) \bar{v}_g^{2n} \|_{\varphi_0}^\# + \| (w_{2n+2} \bar{v}_g^{2n} w_{2n+2}^* - \bar{v}_g^{2n}) \|_{\varphi_0}^\# \\ &\quad + \| (v_g^{2n+2} - 1) (w_{2n+2} \bar{v}_g^{2n} w_{2n+2}^* - \bar{v}_g^{2n}) \|_{\varphi_0}^\#, \end{aligned}$$

we will estimate the above three terms.

Suppose $g \in G_k$. Then for any n with $2n \geq k$, $\varphi_0, \text{Ad} \bar{v}_g^{2n}(\varphi_0) \in \Phi_{2n}^+$, and hence $\|v_g^{2n+2} - 1\|_{\text{Ad} \bar{v}_g^{2n}(\varphi_0)}^\# < 1/4^{2n+2}$ and $\|v_g^{2n+2} - 1\|_{\varphi_0}^\# < 1/4^{2n+2}$ hold by $(3.2n+2)$.

We have

$$\begin{aligned} \|(v_g^{2n+2} - 1) \bar{v}_g^{2n}\|_{\varphi_0}^\# &= \frac{1}{2} (\|(v_g^{2n+2} - 1) \bar{v}_g^{2n}\|_{\varphi_0}^2 + \|\bar{v}_g^{2n*} (v_g^{2n+2*} - 1)\|_{\varphi_0}^2) \\ &= \frac{1}{2} (\|(v_g^{2n+2} - 1)\|_{\text{Ad} \bar{v}_g^{2n}(\varphi_0)}^2 + \|(v_g^{2n+2*} - 1)\|_{\varphi_0}^2) \\ &\leq \|(v_g^{2n+2} - 1)\|_{\text{Ad} \bar{v}_g^{2n}(\varphi_0)}^\# + \|(v_g^{2n+2} - 1)\|_{\varphi_0}^\# \\ &< \frac{2}{16^{2n+2}}. \end{aligned}$$

Hence we get $\|(v_g^{2n+2} - 1) \bar{v}_g^{2n}\|_{\varphi_0}^\# < \sqrt{2}/4^{2n+2} < 1/2^{2n+1}$.

We next estimate $\|w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^\#$. Since $\varphi_0, \bar{v}_g^{2n} \varphi_0, \varphi_0 \bar{v}_g^{2n} \in \Phi'_{2n+1}$, we have $\|[w_{2n+2}, \varphi_0]\| < 1/4^{2n+2}$, $\|[w_{2n+2}, \bar{v}_g^{2n} \varphi_0]\| < 1/4^{2n+2}$ and $\|[w_{2n+2}, \varphi_0 \bar{v}_g^{2n}]\| < 1/4^{2n+2}$ by $(4.2n+2)$. Then

$$\begin{aligned} &\|(w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}) \cdot \varphi_0\| \\ &= \|(w_{2n+2}^* \bar{v}_g^{2n} - \bar{v}_g^{2n} w_{2n+2}^*) w_{2n+2} \cdot \varphi_0\| \\ &\leq \|(w_{2n+2}^* \bar{v}_g^{2n} - \bar{v}_g^{2n} w_{2n+2}^*) \cdot \varphi_0 \cdot w_{2n+2}\| + \|(w_{2n+2}^* \bar{v}_g^{2n} - \bar{v}_g^{2n} w_{2n+2}^*) \cdot [w_{2n+2}, \varphi_0]\| \\ &\leq \|w_{2n+2}^* \bar{v}_g^{2n} \varphi_0 w_{2n+2} - \bar{v}_g^{2n} w_{2n+2}^* \varphi_0 w_{2n+2}\| + 2/4^{2n+2} \\ &\leq \|[w_{2n+2}, \bar{v}_g^{2n} \varphi_0] w_{2n+2}\| + \|\bar{v}_g^{2n} \varphi_0 - \bar{v}_g^{2n} w_{2n+2}^* \varphi_0 w_{2n+2}\| + 2/4^{2n+2} \\ &\leq 1/4^{2n+2} + \|\bar{v}_g^{2n} [\varphi_0, w_{2n+2}^*]\| + 2/4^{2n+2} \\ &\leq 1/4^{2n+1} \end{aligned}$$

holds. Hence

$$\begin{aligned}
\|w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^2 &\leq \|w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}\| \| (w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}) \cdot \varphi_0 \| \\
&\leq 2 \| (w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}) \cdot \varphi_0 \| \\
&\leq \frac{2}{4^{2n+1}}
\end{aligned}$$

holds.

In a similar way, we can show $\|(w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n})^*\|_{\varphi_0}^2 \leq 2/4^{2n+1}$. Hence we get $\|w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^{\#} \leq \sqrt{2/4^{2n+1}} = \sqrt{2}/2^{2n+1} < 1/2^{2n}$.

The third term $\|(v_g^{2n+2} - 1)(w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n})\|_{\varphi_0}^{\#}$ is estimated as follows.

$$\begin{aligned}
\|(v_g^{2n+2} - 1)(w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n})\|_{\varphi_0}^{\#2} &= \frac{1}{2} (\|(v_g^{2n+2} - 1)(w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n})\|_{\varphi_0}^2 \\
&\quad + \|(w_{2n+2}^* \bar{v}_g^{2n*} w_{2n+2} - \bar{v}_g^{2n*})(v_g^{2n+2*} - 1)\|_{\varphi_0}^2) \\
&\leq 2 \|w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^2 + 2 \|v_g^{2n+2*} - 1\|_{\varphi_0}^2 \\
&\leq 4 \|w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^{\#2} + 4 \|v_g^{2n+2*} - 1\|_{\varphi_0}^{\#2} \\
&\leq \frac{4}{4^{2n}} + \frac{4}{4^{2n+2}} \\
&\leq \frac{2}{4^{2n-1}}.
\end{aligned}$$

Hence $\|(v_g^{2n+2} - 1)(w_{2n+2}^* \bar{v}_g^{2n} w_{2n+2} - \bar{v}_g^{2n})\|_{\varphi_0}^{\#} \leq \sqrt{2}/2^{2n-1} < 1/2^{2n-2}$, and we have the following.

$$\begin{aligned}
\|\bar{v}_g^{2n+2} - \bar{v}_g^{2n}\|_{\varphi_0}^{\#} &\leq \frac{1}{2^{2n+1}} + \frac{1}{2^{2n}} + \frac{1}{2^{2n-2}} \\
&\leq \frac{1}{2^{2n-3}}.
\end{aligned}$$

It follows that $\{\bar{v}_g^{2n}\}$ is a Cauchy sequence and converges to some unitary $\hat{v}_g^0$.

In the same way, we can show $\sigma_{2n+1} := \text{Ad} w_{2n+1}^* w_{2n-1}^* \cdots w_1^*$ and $\bar{v}_g^{2n+1}$ converges to $\sigma \in \text{Aut}(M)$ and $\hat{v}_g^1 \in U(M)$ respectively. By (1.n) we get $\text{Ad} \hat{v}_g^0 \circ \theta \circ \alpha_g \circ \theta^{-1} = \text{Ad} \hat{v}_g^1 \circ \sigma \circ \beta_g \circ \sigma^{-1}$, and hence α and β are cocycle conjugate. By the construction, θ, σ are approximately inner.

We will choose a cocycle close to 1. Suppose $\text{Ad} v_g \alpha = \theta \circ \beta_g \circ \theta^{-1}$, $\theta \in \overline{\text{Int}}(M)$. Fix $F \subset\subset G$. Then there exists a unitary w such that $\|w v_g \alpha_g(w^*) - 1\|_{\varphi_0}^{\#} < \varepsilon$ for each $g \in F$. (See the remark after Lemma 6.) Define a new α -cocycle v'_g by $v'_g := w v_g \alpha_g(w^*)$. We then have $\|v'_g - 1\|_{\varphi_0}^{\#} < \varepsilon$ for $g \in F$, and

$$\begin{aligned}
\text{Ad} v'_g \alpha_g &= \text{Ad}(w v_g \alpha_g(w^*)) \circ \alpha_g \\
&= \text{Ad} w \circ \text{Ad} v_g \circ \alpha_g \circ \text{Ad} w^* \\
&= \text{Ad} w \circ \theta \circ \beta_g \circ \theta^{-1} \circ \text{Ad} w^*.
\end{aligned}$$

Put $\sigma := \text{Ad} w \circ \theta$. Then $\sigma \in \overline{\text{Int}}(M)$, and we get $\text{Ad} v'_g \alpha_g = \sigma \circ \beta_g \circ \sigma^{-1}$. $\square$

We present applications of Theorem 3. Let M be an injective factor. By the Connes-Krieger-Haagerup classification of injective factors [2], [12], [5], [7], M is a McDuff factor. Since $\text{Ker mod} = \overline{\text{Int}}(M)$ by [3] and [11], we get the following corollary.

Corollary 7 *Let M be an injective factor, G a discrete amenable group, α, β centrally free actions of G on M . Then $\text{Adv}_g \alpha_g = \theta \circ \beta_g \circ \theta^{-1}$ for some α -cocycle v_g and $\theta \in \overline{\text{Int}}(M)$ if and only if $\text{mod}(\alpha) = \text{mod}(\beta)$. (In the type II_1 case, we regard $\text{mod}(\alpha)$ is trivial for $\alpha \in \text{Aut}(M)$.)*

Theorem 3 can be modified for a relatively McDuff subfactor $N \subset M$ by appropriate changes. Indeed in the proof we only have to replace M_ω with $M_\omega \cap N^\omega$, which is the subfactor version of a central sequence algebra. Especially if $N \subset M$ is a strongly amenable subfactor of type II_1 in the sense of Popa [18], then it is relatively McDuff thanks to Popa's classification theorem of strongly amenable subfactors of type II_1 [18]. We also have $\overline{\text{Int}}(M, N) = \text{Ker } \Phi$ by [13], where $\Phi(\alpha)$ is the Loi invariant for $\alpha \in \text{Aut}(M, N)$, and the equivalence between strong outerness and central freeness by [17]. (Also see [14] for the latter fact.) Hence Theorem 3 gives an alternative proof of the main theorem in [17].

Corollary 8 *Let $N \subset M$ be a strongly amenable subfactor of type II_1 , G a discrete amenable group, and α, β strongly outer actions of G on $N \subset M$. Then $\text{Adv}_g \alpha_g = \theta \circ \beta_g \circ \theta^{-1}$ for some α -cocycle $v_g \in U(N)$ and $\theta \in \overline{\text{Int}}(M, N)$ if and only if $\Phi(\alpha) = \Phi(\beta)$.*

When $N \subset M$ is a strongly amenable subfactor of type II_∞ , then we have $\overline{\text{Int}}(M, N) = \text{Ker } \Phi \cap \text{Ker}(\text{mod})$. Hence we have the following corollary.

Corollary 9 *Let $N \subset M$ be a strongly amenable subfactor of type II_∞ , G a discrete amenable group, and α, β strongly outer actions of G on $N \subset M$. Then $\text{Adv}_g \alpha_g = \theta \circ \beta_g \circ \theta^{-1}$ for some α -cocycle $v_g \in U(N)$ and $\theta \in \overline{\text{Int}}(M, N)$ if and only if $\Phi(\alpha) = \Phi(\beta)$ and $\text{mod}(\alpha) = \text{mod}(\beta)$.*

It is worth noting that Corollary 9 implies the classification of strongly amenable subfactor of type III_λ , $0 < \lambda < 1$, in [13], [17].

References

- [1] Connes, A., *Outer conjugacy classes of automorphisms of factors*, Ann. Sci. Eco. Norm. Sup. **8** (1975), 383–420.
- [2] Connes, A., *Classification of injective factors*, Ann. Math. **104** (1976), 73–115.
- [3] Connes, A., *On the classification of von Neumann algebras and their automorphisms*, Symp. Math. **XX** (1976), 435–478.
- [4] Connes, A., *Periodic automorphisms of the hyperfinite factor of type II_1* , Acta Sci. Math **39** (1977), 39–66.
- [5] Connes, A., *Type III_1 factors, property L'_λ and closure of inner automorphisms*, J. Operator Theory **14** (1985), 189–211.
- [6] Evans, D. E. and Kishimoto, A., *Trace scaling automorphisms of certain stable AF algebras*, Hokkaido Math J. **26** (1997), 211–224.

- [7] Haagerup, U., *Connes' bicentralizer problem and uniqueness of the injective factor of type III_1* , Acta Math. **158** (1987), 95–148.
- [8] Izumi, M., *Finite group actions on C^* -algebras with the Rohlin property I*, Duke Math. J. **122** (2004), 233–280.
- [9] Jones, V. F. R., *Action of finite groups on the hyperfinite type II_1 factor*, Memoirs of Amer. Math. Soc **237** (1980).
- [10] Katayama, Y., Sutherland, C. E., and Takesaki, M., *The characteristic square of a factor and the cocycle conjugacy of discrete group actions on factors*, Invent. Math. **132** (1998), 331–380.
- [11] Kawahigashi, Y., Sutherland, C. E., and Takesaki, M., *The structure of the automorphism group of an injective factor and the cocycle conjugacy of discrete abelian group actions*, Acta Math. **169** (1992), 105–130.
- [12] Krieger, W., *On ergodic flows and the isomorphism of factors*, Math. Annal. **223** (1976), 19–70.
- [13] Loi, P., *On automorphisms of subfactors*, J. Funct. Anal. **141** (1996), 275–293.
- [14] Masuda, T., *An analogue of Connes-Haagerup approach to classification of subfactors of type III_1* , preprint.
- [15] Nakamura, H., *Aperiodic automorphisms of nuclear purely infinite simple C^* -algebras*, Ergod. Th. and Dynam. Sys. **20** (2000), 1749–1765.
- [16] Ocneanu, A., *Action of discrete amenable groups on von Neumann algebras*, vol. 1138, Springer, Berlin, (1985).
- [17] Popa, S., *Classification actions of discrete amenable groups on subfactors of type II*, preprint, (1992).
- [18] Popa, S., *Classification of amenable subfactor of type II*, Acta Math. **172** (1994), 163–255.
- [19] Sutherland, C. E. and Takesaki, M., *Actions of discrete amenable groups on injective factors of type III_λ , $\lambda \neq 1$* , Pacific. J. Math. **137** (1989), 405–444.